



Vegetable rumen protected fat



- Lipitec Bovi LM is a sprayed fat, developed especially for increased production of milk and butterfat in high yielding dairy cows
- Improves fat content in milk
- Extra milk
- Lipitec Bovi LM can be fed in pure form or used in compound feed
- Lipitec Bovi LM is based on saturated fatty acids, enriched with mono/diglycerides, to maximize digestibility
- VLOG verified

Typical analyses:

i ypicai alialyses.	
Crude fat:	Min. 99 %
Moisture and impurities:	Max. 1 %
NEL:	24 MJ/kg
ME:	36 MJ/kg
Fatty acids, % of crude fat:	90 %
lodine value:	Max. 10
Melting point:	57° C
Ni content:	<5 ppm
Colour:	Light brown (content of carotene)

Fatty acid profile:

C _{14:0} (myristic acid):	Max. 1,5 %
C _{16:0} (palmetic acid):	40-55 %
C _{18:0} (stearic acid).	40-55 %
C _{18:1} (oleic acid):	Max. 8 %
Others:	Max. 3 %

Recommended use:

Lipitec Bovi LM can be fed directly upon silage or other feed.		
Max. recommended daily supply for dairy cows:		
	•	
0-20 weeks post calving:	500 gram	
>20 weeks post calving:	250 gram	
, , , , ,		

Can be delivered in small bags, big bags, bulk or liquid. Bags are recommended to be closed between use, since mono glycerides can absorb moisture from humid environment. Keep dry and out of direct sunlight. Expiry date: 1 year from production.





CVR-nr. DK13408785 – PVR-nr. α-208-G756968 www.lipitec.com - info@lipitec.com

